

# Degreeewise $n$ -projective and $n$ -flat model structures on chain complexes

Marco A. Pérez B.

Université du Québec à Montréal.

Département de Mathématiques.

July 1st, 2012

## Abstract

In [4], the authors construct an abelian model structure on the category of chain complexes over left  $R$ -modules,  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ , where the class of (trivially) cofibrant objects is given by the class of degreeewise projective (resp. exact) chain complexes. Using a generalization of a well known theorem by I. Kaplansky, we generalize the method used in [4] in order to obtain, for each integer  $n > 0$ , a new abelian model structure on the  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ , where the class of (trivially) cofibrant objects is the class of (resp. exact) chain complexes whose terms have projective dimension at most  $n$ , provided the ring  $R$  is right noetherian. We also give another method to construct this model structure, which also works to construct a model structure where the class of (trivially) cofibrant objects is given by the class of (resp. exact) chain complexes whose terms have flat dimension at most  $n$ .

## 1 Introduction

A cotorsion pair in an abelian category  $\mathcal{C}$  is a pair  $(\mathcal{A}, \mathcal{B})$ , where  $\mathcal{A}$  and  $\mathcal{B}$  are classes of objects of  $\mathcal{C}$  such that they are orthogonal to each other with respect to the Ext functor. A model category is a bicomplete category with three classes of morphisms, called cofibrations, fibrations and weak equivalences, satisfying certain conditions. It turns out to be that these two notions have a deep connection. As far as the author knows, the first person who described this connection was M. Hovey in the paper *Cotorsion pairs, model category structures and representation theory*, where he proved

that any two compatible and complete cotorsion pairs  $(\mathcal{A}, \mathcal{B} \cap \mathcal{E})$  and  $(\mathcal{A} \cap \mathcal{E}, \mathcal{B})$ , in a bicomplete abelian category  $\mathcal{C}$ , give rise to a unique abelian model structure on  $\mathcal{C}$  where  $\mathcal{A}$  is the class of cofibrant objects,  $\mathcal{B}$  is the class of fibrant objects, and  $\mathcal{E}$  is the class of trivial objects.

From this point there has been an increasing interest in constructing new model structures, specially on  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ . One of the most influential researchers in this matter has been J. Gillespie, who has provided several results that allows us to induce cotorsion pairs in the category  $\mathbf{Ch}(\mathcal{C})$  of chain complexes over an abelian category  $\mathcal{C}$ , from a certain cotorsion pair in  $\mathcal{C}$ . One of those results states that given a cotorsion pair  $(\mathcal{A}, \mathcal{B})$  in an abelian category  $\mathcal{C}$  with enough projective and injective objects, there exist two cotorsion pairs in  $\mathbf{Ch}(\mathcal{C})$  given by  $(\mathrm{dw}\widetilde{\mathcal{A}}, (\mathrm{dw}\widetilde{\mathcal{A}})^\perp)$  and  $(\mathrm{ex}\widetilde{\mathcal{A}}, (\mathrm{ex}\widetilde{\mathcal{A}})^\perp)$ , where  $\mathrm{dw}\widetilde{\mathcal{A}}$  is the class of chain complexes  $X$  such that  $X_m \in \mathcal{A}$  for every  $m \in \mathbb{Z}$ , and  $\mathrm{ex}\widetilde{\mathcal{A}} = \mathrm{dw}\widetilde{\mathcal{A}} \cap \mathcal{E}$  where  $\mathcal{E}$  is the class of exact complexes. As an example, if  $\mathcal{P}_0$  denotes the class of projective modules in the category  $R\text{-}\mathbf{Mod}$  of left  $R$ -modules, then the cotorsion pair  $(\mathcal{P}_0, R\text{-}\mathbf{Mod})$  induces two cotorsion pairs  $(\mathrm{dw}\widetilde{\mathcal{P}_0}, (\mathrm{dw}\widetilde{\mathcal{P}_0})^\perp)$  and  $(\mathrm{ex}\widetilde{\mathcal{P}_0}, (\mathrm{ex}\widetilde{\mathcal{P}_0})^\perp)$ . In [4], the authors prove that these pairs are compatible and complete, with the help of a theorem by I. Kaplansky, namely that every projective module can be written as a direct sum of countably generated projective modules. Then, using [10, Theorem 2.2], they get a new abelian model structure on  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$  where the class of cofibrant objects is the class  $\mathrm{dw}\widetilde{\mathcal{P}_0}$ , which we shall call the class of **degree-wise projective complexes**. We shall refer to this model structure as the **dw-projective model structure**.

In [2] it is proven that if  $\mathcal{P}_n$  denotes the class of left  $R$ -modules with projective dimension at most  $n$ , then  $(\mathcal{P}_n, \mathcal{P}_n^\perp)$  is a complete and hereditary cotorsion pair. It follows we have two induced cotorsion pairs  $(\mathrm{dw}\widetilde{\mathcal{P}_n}, (\mathrm{dw}\widetilde{\mathcal{P}_n})^\perp)$  and  $(\mathrm{ex}\widetilde{\mathcal{P}_n}, (\mathrm{ex}\widetilde{\mathcal{P}_n})^\perp)$  in  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ . Our goal is to prove that these two cotorsion pairs are complete for

every  $n > 0$ , in order to obtain a new abelian model structure on  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$  such that  $\text{dw}\widetilde{\mathcal{P}}_n$  is the class of cofibrant objects.

This paper is organized as follows. First, we recall some definitions and introduce the notation we shall use. Then, we shall give a “generalization” of the Kaplansky Theorem in  $\text{Mod-}R$  provided that  $R$  is left noetherian. Specifically, we shall prove that every module of projective dimension  $\leq n$  has a  $\mathcal{P}_n^{\aleph_0}$ -filtration, where  $\mathcal{P}_n^{\aleph_0}$  is the set of all modules  $M$  for which there exists an exact sequence

$$0 \longrightarrow P_n \longrightarrow \cdots \longrightarrow P_1 \longrightarrow P_0 \longrightarrow M \longrightarrow 0$$

where  $P_k$  is a countably generated projective module, for every  $0 \leq k \leq n$ . Using this result, we shall prove that  $(\text{dw}\widetilde{\mathcal{P}}_n, (\text{dw}\widetilde{\mathcal{P}}_n)^\perp)$  and  $(\text{ex}\widetilde{\mathcal{P}}_n, (\text{ex}\widetilde{\mathcal{P}}_n)^\perp)$  are complete cotorsion pairs. Then, we shall give another method to prove the previous result. The interesting thing of this other method, based on arguments appearing in the proof of [1, Proposition 4.1], is that it can be applied to show that  $(\text{dw}\widetilde{\mathcal{F}}_n, (\text{dw}\widetilde{\mathcal{F}}_n)^\perp)$  and  $(\text{ex}\widetilde{\mathcal{F}}_n, (\text{ex}\widetilde{\mathcal{F}}_n)^\perp)$  are complete cotorsion pairs, where  $\mathcal{F}_n$  denotes the class of left  $R$ -modules having flat dimension at most  $n$ . At the end of the paper, we shall give some comments concerning the  $\text{dw-}n$ -projective and  $\text{dw-}n$ -flat model structures.

## 2 Preliminaries

This section is devoted to recall some notions and to introduce part of the notation we shall use throughout the paper. From now on, we work in the category  $R\text{-}\mathbf{Mod}$  of left  $R$ -modules, and the category  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$  of chain complexes over  $R\text{-}\mathbf{Mod}$ . Given a chain complex  $X = (X_m)_{m \in \mathbb{Z}}$  with boundary maps  $\partial_m^X : X_m \longrightarrow X_{m-1}$ , we shall denote  $Z_m(X) := \text{Ker}(\partial_m^X)$ . A chain complex  $X$  is said to be **exact** if  $Z_m(X) = \partial_{m+1}(X_{m+1})$ , for every  $m \in \mathbb{Z}$ . A chain complex  $Y$  is said to be a **subcomplex** of  $X$  if there exists a monomorphism  $i : Y \longrightarrow X$ . Then we can define the **quotient**

**complex**  $X/Y$  as the complex whose components are given by  $(X/Y)_m = X_m/Y_m$  and whose boundary maps  $\partial_m^{X/Y} : X_m/Y_m \longrightarrow X_{m-1}/Y_{m-1}$  are given by

$$x + Y_m \mapsto \partial_m^X(x) + Y_{m-1}.$$

Let  $\mathcal{C}$  be either  $R\text{-}\mathbf{Mod}$  or  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ . Let  $\mathcal{A}$  and  $\mathcal{B}$  be two classes of objects in  $\mathcal{C}$ . The pair  $(\mathcal{A}, \mathcal{B})$  is called a **cotorsion pair** in  $\mathcal{C}$  if the following conditions are satisfied:

- (1)  $\mathcal{A} = {}^\perp \mathcal{B} := \{X \in \text{Ob}(\mathcal{C}) \mid \text{Ext}^1(X, B) = 0 \text{ for every } B \in \mathcal{B}\}.$
- (2)  $\mathcal{B} = \mathcal{A}^\perp := \{X \in \text{Ob}(\mathcal{C}) \mid \text{Ext}^1(A, X) = 0 \text{ for every } A \in \mathcal{A}\}.$

A cotorsion pair  $(\mathcal{A}, \mathcal{B})$  in  $\mathcal{C}$  is said to be **complete** if:

- (a)  $(\mathcal{A}, \mathcal{B})$  has **enough projectives**: for every object  $X$  there exist objects  $A \in \mathcal{A}$  and  $B \in \mathcal{B}$ , and a short exact sequence

$$0 \longrightarrow B \longrightarrow A \longrightarrow X \longrightarrow 0.$$

- (b)  $(\mathcal{A}, \mathcal{B})$  has **enough injectives**: for every object  $X$  there exist objects  $A' \in \mathcal{A}$  and  $B' \in \mathcal{B}$ , and a short exact sequence

$$0 \longrightarrow X \longrightarrow B' \longrightarrow A' \longrightarrow 0.$$

A cotorsion pair  $(\mathcal{A}, \mathcal{B})$  is said to be **cogenerated** by a set  $\mathcal{S} \subseteq \mathcal{A}$  if  $\mathcal{B} = \mathcal{S}^\perp$ . There is a wide range of complete cotorsion pairs, thanks to the following result, known as the Eklof and Trlifaj Theorem.

**Theorem 2.1.** [5, Theorem 10] *Every cotorsion pair in  $\mathcal{C}$  cogenerated by a set is complete.*

**Example 2.1.**

- (1) If  $\mathcal{P}_0$  denotes the class of projective modules, then  $(\mathcal{P}_0, R\text{-}\mathbf{Mod})$  is a cotorsion pair. Since every projective module is a direct summand of a free module, and  $R$  is projective, one can show that  $(\mathcal{P}_0, R\text{-}\mathbf{Mod})$  is cogenerated by the set  $\{R\}$  and hence it is complete.
- (2) Similarly, if  $\mathcal{I}_0$  denotes the class of injective modules, then  $(R\text{-}\mathbf{Mod}, \mathcal{I}_0)$  is a cotorsion pair. Using the Baer's Criterion, one can show that  $(R\text{-}\mathbf{Mod}, \mathcal{I}_0)$  is cogenerated by the set of modules of the form  $R/I$ , where  $I$  is a left ideal of  $R$ . So  $(R\text{-}\mathbf{Mod}, \mathcal{I}_0)$  is a complete cotorsion pair.
- (3) A less trivial example of a complete cotorsion pair is given by the **flat cotorsion pair**  $(\mathcal{F}_0, \mathcal{F}_0^\perp)$ , where  $\mathcal{F}_0$  is the class of flat modules. This result was proven by Edgard E. Enochs by using the Eklof and Trlifaj Theorem. Enochs proved that the pair  $(\mathcal{F}_0, \mathcal{F}_0^\perp)$  is cogenerated by the set  $\mathcal{S} = \{S \in \mathcal{F}_0 : \text{Card}(S) \leq \kappa\}$ , where  $\kappa$  is an infinite cardinal with  $\kappa \geq \text{Card}(R)$ .
- (4) The following example is probably the most important cotorsion pair we shall consider in this paper, the pair  $(\mathcal{P}_n, \mathcal{P}_n^\perp)$ , where  $\mathcal{P}_n$  is the class of modules which have projective dimension  $\leq n$ . Recall that a module  $M$  has **projective dimension**  $\leq n$  if there exists an exact sequence

$$0 \longrightarrow P_n \longrightarrow P_{n-1} \longrightarrow \cdots \longrightarrow P_1 \longrightarrow P_0 \longrightarrow M \longrightarrow 0,$$

such that  $P_k$  is a projective module, for every  $0 \leq k \leq n$ . Such a sequence is called a **projective resolution of  $M$  of length  $n$** . We shall refer to the modules in  $\mathcal{P}_n$  as  **$n$ -projective modules**. In [2], the authors proved that  $(\mathcal{P}_n, \mathcal{P}_n^\perp)$  is a cotorsion pair cogenerated by the set of all  $n$ -projective modules whose cardinality is less or equal than a given infinite cardinal  $\kappa$  with  $\kappa \geq \text{Card}(R)$ .

(5) In a similar way, consider the class  $\mathcal{F}_n$  of modules  $M$  such that  $M$  has flat dimension at most  $n$ , or equivalently, there is an exact sequence

$$0 \longrightarrow F_n \longrightarrow \cdots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow M \longrightarrow 0$$

where  $F_k$  is a flat module, for every  $0 \leq k \leq n$ . This sequence is called a **flat resolution of length  $n$** . In [9, Theorem 4.1.3], it is proven that  $(\mathcal{F}_n, \mathcal{F}_n^\perp)$  is a complete cotorsion pair. Here we shall give a easier proof of this fact.

Now we recall the notion of a model category. Given a category  $\mathcal{C}$ , a map  $f$  in  $\mathcal{C}$  is a **retract** of a map  $g$  in  $\mathcal{C}$  if there is a commutative diagram of the form

$$\begin{array}{ccccc} A & \longrightarrow & C & \longrightarrow & A \\ f \downarrow & & g \downarrow & & \downarrow f \\ B & \longrightarrow & D & \longrightarrow & B \end{array}$$

where the horizontal composites are identities. Let  $f : A \longrightarrow B$  and  $g : C \longrightarrow D$  be two maps in  $\mathcal{C}$ . We shall say that  $f$  has the **left lifting property** with respect to  $g$  (or that  $g$  has the **right lifting property** with respect to  $f$ ) if for every pair of maps  $u : A \longrightarrow C$  and  $v : B \longrightarrow D$  with  $g \circ u = v \circ f$ , there exists a map  $d : B \longrightarrow C$  such that  $g \circ d = v$  and  $d \circ f = u$ .

$$\begin{array}{ccc} \begin{array}{ccc} A & \xrightarrow{u} & C \\ f \downarrow & & \downarrow g \\ B & \xrightarrow{v} & D \end{array} & \rightsquigarrow & \begin{array}{ccc} A & \xrightarrow{u} & C \\ f \downarrow & \exists d \nearrow & \downarrow g \\ B & \xrightarrow{v} & D \end{array} \end{array}$$

A **model category** is a bicomplete category  $\mathcal{C}$  equipped with three classes of maps named **cofibrations**, **fibrations** and **weak equivalences**, satisfying the following properties:

- (1) **3 for 2:** If  $f$  and  $g$  are maps of  $\mathcal{C}$  such that  $g \circ f$  is defined and two of  $f$ ,  $g$  and  $g \circ f$  are weak equivalences, then so is the third.
- (2) If  $f$  and  $g$  are maps of  $\mathcal{C}$  such that  $f$  is a retract of  $g$  and  $g$  is a weak equivalence, cofibration, or fibration, then so is  $f$ .

Define a map to be a **trivial cofibration** if it is both a weak equivalence and a cofibration. Similarly, define a map to be a **trivial fibration** if it is both a weak equivalence and a fibration.

- (3) Trivial cofibrations have the left lifting property with respect to fibrations, and cofibrations have the left lifting property with respect to trivial fibrations.
- (4) Every map  $f$  can be factored as  $f = \alpha \circ \beta = \gamma \circ \delta$ , where  $\alpha$  (resp.  $\delta$ ) is a cofibration (resp. fibration), and  $\gamma$  (resp.  $\beta$ ) is a trivial cofibration (resp. trivial fibration).

An object  $X$  in  $\mathcal{C}$  is called **cofibrant** if the map  $0 \rightarrow X$  is a cofibration, **fibrant** if the map  $X \rightarrow 1$  is a fibration, and **trivial** if the map  $0 \rightarrow X$  is a weak equivalence, where  $0$  and  $1$  denote the initial and terminal objects of  $\mathcal{C}$ , respectively.

Given a bicomplete abelian category  $\mathcal{C}$ , a model structure on it is said to be **abelian** if the following conditions are satisfied:

- (a) A map is a cofibration if and only if it is a monomorphism with cofibrant cokernel.
- (b) A map is a fibration if and only if it is an epimorphism with fibrant kernel.

### 3 Degreewise $n$ -projective complexes

We begin this section with the notion of a filtration. Let  $\mathcal{C}$  be either  $R\text{-}\mathbf{Mod}$  or  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ . Given an object  $X \in \mathcal{C}$ , by a **filtration** of  $X$  indexed by an ordinal  $\lambda$  we shall mean a family  $(X^\alpha : \alpha < \lambda)$  of subobjects of  $X$  such that:

- (1)  $X = \bigcup_{\alpha < \lambda} X_\alpha$ .
- (3)  $X^\alpha$  is a subobject of  $X^{\alpha'}$  whenever  $\alpha \leq \alpha'$ .
- (4)  $X^\beta = \bigcup_{\alpha < \beta} X^\alpha$  for any limit ordinal  $\beta < \lambda$ .

If  $\mathcal{S}$  is some class of objects in  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ , we say that a filtration  $(X^\alpha : \alpha < \lambda)$  of  $X$  is a  **$\mathcal{S}$ -filtration** if for each  $\alpha + 1 < \lambda$  we have that  $X_0$  and  $X^{\alpha+1}/X^\alpha$  are isomorphic to an element of  $\mathcal{S}$ .

The construction of the model structure given in [4] is based on a theorem by I. Kaplansky, namely:

**Theorem 3.1** (Kaplansky's Theorem). *If  $P$  is a projective module then  $P$  is a direct sum of countable generated projective modules.*

So when one thinks of a possible generalization of the dw-projective model structure for  $n$ -projective modules, a good question would be if it is possible to generalize the Kaplansky's Theorem for such modules. Let  $M \in \mathcal{P}_n$  be an  $n$ -projective module:

$$0 \longrightarrow P_n \longrightarrow \cdots \longrightarrow P_1 \longrightarrow P_0 \longrightarrow M \longrightarrow 0.$$

By Kaplansky's Theorem we can write  $P_k = \bigoplus_{i \in I_k} P_k^i$ , where  $P_k^i$  is a countably generated projective module, for every  $i \in I_k$  and every  $0 \leq k \leq n$ . Then we can rewrite the previous resolution as

$$0 \longrightarrow \bigoplus_{i \in I_n} P_n^i \longrightarrow \bigoplus_{i \in I_{n-1}} P_{n-1}^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1} P_1^i \longrightarrow \bigoplus_{i \in I_0} P_0^i \longrightarrow M \longrightarrow 0.$$

From now on we shall write any projective resolution of length  $n$  by using such direct sum decompositions. We shall denote by  $\mathcal{P}_n^{\aleph_0}$  the set of all modules  $M$  having a projective resolution as above, where  $I_k$  is a countable set for each  $0 \leq k \leq n$ .

For any class of modules  $\mathcal{A}$ , we denote by  $\text{dw}\widetilde{\mathcal{A}}$  (resp.  $\text{ex}\widetilde{\mathcal{A}}$ ) the class of (resp. exact) chain complexes such that each term belongs to  $\mathcal{A}$ . We shall prove that  $(\text{dw}\widetilde{\mathcal{P}}_n, (\text{dw}\widetilde{\mathcal{P}}_n)^\perp)$  is a cotorsion pair cogenerated by the set  $\text{dw}\widetilde{\mathcal{P}}_n^{\aleph_0}$ . We shall name  $\text{dw}\widetilde{\mathcal{P}}_n$  the class of **dw- $n$ -projective chain complexes**. The fact that  $(\text{dw}\widetilde{\mathcal{P}}_n, (\text{dw}\widetilde{\mathcal{P}}_n)^\perp)$  is a cotorsion pair in  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$  is a consequence of the following result (which is proven by its author for any abelian category):

**Proposition 3.1.** [8, Proposition 3.2] *Let  $(\mathcal{A}, \mathcal{B})$  be a cotorsion pair in  $R\text{-}\mathbf{Mod}$ . Then  $(\text{dw}\widetilde{\mathcal{A}}, (\text{dw}\widetilde{\mathcal{A}})^\perp)$  is a cotorsion pair in  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ .*

We shall prove that every dw- $n$ -projective complex has a  $\text{dw}\widetilde{\mathcal{P}}_n^{\aleph_0}$ -filtration. Then the completeness of  $(\text{dw}\widetilde{\mathcal{P}}_n, (\text{dw}\widetilde{\mathcal{P}}_n)^\perp)$  shall be a consequence of Theorem 2.1 and the following result:

**Proposition 3.2.** *Let  $(\mathcal{A}, \mathcal{B})$  be a cotorsion pair in  $\mathcal{C} = R\text{-}\mathbf{Mod}, \mathbf{Ch}(R\text{-}\mathbf{Mod})$  and let  $\mathcal{S} \subseteq \mathcal{A}$  be a set of objects of  $\mathcal{C}$ . If every  $A \in \mathcal{A}$  has a  $\mathcal{S}$ -filtration, then  $(\mathcal{A}, \mathcal{B})$  is cogenerated by  $\mathcal{S}$ .*

Before proving this, we need the following result known as the Eklof's Lemma. For a proof of this we refer the reader to [9, Lemma 3.1.2] or [6, Theorem 7.3.4].

**Lemma 3.1** (Eklof's Lemma). *In  $\mathcal{C} = R\text{-}\mathbf{Mod}, \mathbf{Ch}(R\text{-}\mathbf{Mod})$  let  $A$  and  $B$  be two objects. If  $A$  has a  ${}^\perp\{B\}$ -filtration, then  $A \in {}^\perp\{B\}$ .*

**Proof of Proposition 3.2:** Consider the cotorsion pair  $({}^\perp(\mathcal{S}^\perp), \mathcal{S}^\perp)$ . We shall show that  $({}^\perp(\mathcal{S}^\perp), \mathcal{S}^\perp) = (\mathcal{A}, \mathcal{B})$ . It suffices to show that  $\mathcal{B} = \mathcal{S}^\perp$ , since this equality implies  $\mathcal{A} = {}^\perp\mathcal{B} = {}^\perp(\mathcal{S}^\perp)$ . Since  $\mathcal{S} \subseteq \mathcal{A}$ , we have  $\mathcal{B} = \mathcal{A}^\perp \subseteq \mathcal{S}^\perp$ . Now let  $Y \in \mathcal{S}^\perp$ ,  $A \in \mathcal{A}$  and

let  $(A_\alpha : \alpha < \lambda)$  be an  $\mathcal{S}$ -filtration of  $A$ . We have

$$\begin{aligned}\mathrm{Ext}^1(A_0, Y) &= \mathrm{Ext}^1(0, Y) = 0, \\ \mathrm{Ext}^1(A_{\alpha+1}/A_\alpha, Y) &=, \text{ whenever } \alpha + 1 < \lambda,\end{aligned}$$

since  $A_0$  and  $A_{\alpha+1}/A_\alpha$  are isomorphic to objects in  $\mathcal{C}$ . Then  $(A_\alpha : \alpha < \lambda)$  is a  ${}^\perp\{Y\}$ -filtration of  $A$ . By the Eklof's Lemma, we have  $\mathrm{Ext}^1(A, Y) = 0$ , i.e.  $Y \in \mathcal{A}^\perp = \mathcal{B}$  since  $A$  is any module in  $\mathcal{A}$ . Hence  $\mathcal{S}^\perp \subseteq \mathcal{B}$ .  $\square$

In order to construct  $\widetilde{\mathrm{dw}\mathcal{P}_n^{\aleph_0}}$ -filtrations of dw- $n$ -projective complexes, we need the following generalization of the Kaplansky's Theorem:

**Lemma 3.2** (Kaplansky's Theorem for  $n$ -projective modules). *Let  $R$  be a noetherian ring. Let  $M \in \mathcal{P}_n$  and let  $N$  be a countably generated submodule of  $M$ . Then there exists a  $\mathcal{P}_n^{\aleph_0}$ -filtration of  $M$ , say  $(M_\alpha : \alpha < \lambda)$  with  $\lambda > 1$ , such that  $M_1 \in \mathcal{P}_n^{\aleph_0}$  and  $N \subseteq M_1$ .*

*Proof.* Let  $M \in \mathcal{P}_n$  and let

$$0 \longrightarrow \bigoplus_{i \in I_n} P_n^i \longrightarrow \bigoplus_{i \in I_{n-1}} P_{n-1}^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1} P_1^i \longrightarrow \bigoplus_{i \in I_0} P_0^i \longrightarrow M \longrightarrow 0$$

be a projective resolution of  $M$ . We shall construct a  $\mathcal{P}_n^{\aleph_0}$ -filtration  $(M_\alpha : \alpha < \lambda)$  of  $M$ , with  $N \subseteq M_1$ , by using transfinite induction. For  $\alpha = 0$  set  $M_0 = 0$ . Now we construct  $M_1$ . Let  $\mathcal{G}$  be a countable set of generators of  $N$ . Since  $f_0$  is surjective, for every  $g \in \mathcal{G}$  we can choose  $y_g \in \bigoplus_{i \in I_0} P_0^i$  such that  $g = f_0(y_g)$ . Consider the set  $Y = \{y_g : g \in \mathcal{G}\}$ . Since  $Y$  is a countable subset of  $\bigoplus_{i \in I_0} P_0^i$ , we have that  $\langle Y \rangle$  is a countably generated submodule of  $P_0$ . Choose a countable subset  $I_0^{1,0} \subseteq I_0$  such that  $\langle Y \rangle \subseteq \bigoplus_{i \in I_0^{1,0}} P_0^i$ . Then  $f_0(\langle Y \rangle) \subseteq N$ . Consider  $\mathrm{Ker} \left( f_0|_{\bigoplus_{i \in I_0^{1,0}} P_0^i} \right)$ . Since  $\bigoplus_{i \in I_0^{1,0}} P_0^i$  is countably generated and  $\mathrm{Ker} \left( f_0|_{\bigoplus_{i \in I_0^{1,0}} P_0^i} \right)$  is a submodule of

$\bigoplus_{i \in I_0^{1,0}} P_0^i$ , we have that  $\text{Ker} \left( f_0|_{\bigoplus_{i \in I_0^{1,0}} P_0^i} \right)$  is also countably generated, since  $R$  is noetherian. Let  $\mathcal{B}$  be a countable set of generators of  $\text{Ker} \left( f_0|_{\bigoplus_{i \in I_0^{1,0}} P_0^i} \right)$ . Let  $b \in \mathcal{B}$ , then  $f(b) = 0$  and by exactness of the above sequence there exists  $y_b \in \bigoplus_{i \in I_1} P_1^i$  such that  $b = f_1(y_b)$ . Let  $Y' = \{y_b : b \in \mathcal{B}\}$ . Note that  $Y'$  is a countable subset of  $(f_1)^{-1} \left( \text{Ker} \left( f_0|_{\bigoplus_{i \in I_0^{1,0}} P_0^i} \right) \right)$ . Then  $\langle Y' \rangle$  is a countably generated submodule of  $\bigoplus_{i \in I_1} P_1^i$ . Hence there exists a countable subset  $I_1^{1,0} \subseteq I_1$  such that  $\bigoplus_{i \in I_1^{1,0}} P_1^i \supseteq \langle Y' \rangle$ . Thus  $f_1 \left( \bigoplus_{i \in I_1^{1,0}} P_1^i \right) \supseteq f_1(\langle Y' \rangle)$ . Now let  $z \in \text{Ker} \left( f_0|_{\bigoplus_{i \in I_0^{1,0}} P_0^i} \right)$ . Then  $z = r_1 b_1 + \dots + r_m b_m$ , where each  $b_j \in \mathcal{B}$ . Since  $b_j = f_1(y_{b_j})$  with  $y_{b_j} \in Y'$ , we get  $z = f_1(r_1 y_{b_1} + \dots + r_m y_{b_m}) \in f_1(\langle Y' \rangle)$ . Hence,  $\text{Ker} \left( f_0|_{\bigoplus_{i \in I_0^{1,0}} P_0^i} \right) \subseteq f_1(\langle Y' \rangle) \subseteq f_1 \left( \bigoplus_{i \in I_1^{1,0}} P_1^i \right)$ . Use the same argument to find a countable subset  $I_2^{1,0} \subseteq I_2$  such that  $f_2 \left( \bigoplus_{i \in I_2^{1,0}} P_2^i \right) \supseteq \text{Ker} \left( f_1|_{\bigoplus_{i \in I_1^{1,0}} P_1^i} \right)$ . Repeat the same argument until find a countable subset  $I_n^{1,0} \subseteq I_n$  such that  $f_n \left( \bigoplus_{i \in I_n^{1,0}} P_n^i \right) \supseteq \text{Ker} \left( f_{n-1}|_{\bigoplus_{i \in I_{n-1}^{1,0}} P_{n-1}^i} \right)$ . Now,  $f_n \left( \bigoplus_{i \in I_n^{1,0}} P_n^i \right)$  is a countably generated submodule of  $\bigoplus_{i \in I_{n-1}} P_{n-1}^i$ . Then choose a countable subset  $I_{n-1}^{1,0} \subseteq I_{n-1}^{1,1} \subseteq I_{n-1}$  such that  $f_n \left( \bigoplus_{i \in I_n^{1,0}} P_n^i \right) \subseteq \bigoplus_{i \in I_{n-1}^{1,1}} P_{n-1}^i$ . Repeat this process until find a countable subset  $I_0^{1,0} \subseteq I_0^{1,1} \subseteq I_0$  satisfying  $f_1 \left( \bigoplus_{i \in I_1^{1,1}} P_1^i \right) \subseteq \bigoplus_{i \in I_0^{1,1}} P_0^i$ . Now choose a countable subset  $I_1^{1,1} \subseteq I_1^{1,2} \subseteq I_1$  such that  $f_1 \left( \bigoplus_{i \in I_2^{1,2}} P_2^i \right) \supseteq \text{Ker} \left( f_0|_{\bigoplus_{i \in I_0^{1,1}} P_0^i} \right)$ . What we have been doing so far is called the zig-zag procedure. Keep repeating this procedure infinitely many times, and set  $I_k^1 = \bigcup_{m \geq 0} I_k^{1,m}$ , for every  $0 \leq k \leq n$ . By construction, we get the following exact sequence

$$0 \longrightarrow \bigoplus_{i \in I_n^1} P_n^i \longrightarrow \bigoplus_{i \in I_{n-1}^1} P_{n-1}^i \longrightarrow \dots \longrightarrow \bigoplus_{i \in I_1^1} P_1^i \longrightarrow \bigoplus_{i \in I_0^1} P_0^i \longrightarrow M_1 \longrightarrow 0$$

where  $x \in M_1 := \text{CoKer} \left( \bigoplus_{i \in I_1^1} P_1^i \longrightarrow \bigoplus_{i \in I_0^1} P_0^i \right) \subseteq M$  and  $N \subseteq M_1$ . We take the quotient of the resolution of  $M$  by the resolution of  $M'$ , and get the following commutative diagram:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \bigoplus_{i \in I_n^1} P_n^i & \longrightarrow & \cdots & \longrightarrow & \bigoplus_{i \in I_1^1} P_1^i & \longrightarrow & \bigoplus_{i \in I_0^1} P_0^i & \longrightarrow & M_1 & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \bigoplus_{i \in I_n} P_n^i & \longrightarrow & \cdots & \longrightarrow & \bigoplus_{i \in I_1} P_1^i & \longrightarrow & \bigoplus_{i \in I_0} P_0^i & \longrightarrow & M & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \bigoplus_{i \in I_n - I_n^1} P_n^i & \longrightarrow & \cdots & \longrightarrow & \bigoplus_{i \in I_1 - I_1^1} P_1^i & \longrightarrow & \bigoplus_{i \in I_0 - I_0^1} P_0^i & \longrightarrow & M/M_1 & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
& & 0 & & 0 & & 0 & & 0 & & 0 & & 
\end{array}$$

where the third row is an exact sequence since the class of exact complexes is closed under taking cokernels. Then we have a projective resolution of length  $n$  for  $M/M_1$ . Repeat the same procedure above for  $M/M_1$ , by choosing  $x^1 + M_1 \in M/M_1 - \{0 + M_1\}$ , in order to get an exact sequence

$$0 \longrightarrow \bigoplus_{i \in I_n^2 - I_n^1} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^2 - I_1^1} P_1^i \longrightarrow \bigoplus_{i \in I_0^2 - I_0^1} P_0^i \longrightarrow M_2/M_1 \longrightarrow 0,$$

for some module  $M_1 \subseteq M_2 \subseteq M$ , such that  $I_k^2 - I_k^1$  is countable for every  $0 \leq k \leq n$ . Note that

$$0 \longrightarrow \bigoplus_{i \in I_n^2} P_n^i \longrightarrow \bigoplus_{i \in I_{n-1}^2} P_{n-1}^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^2} P_1^i \longrightarrow \bigoplus_{i \in I_0^2} P_0^i \longrightarrow M_2 \longrightarrow 0$$

is a projective resolution of  $M_2$ , since we have a commutative diagram

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \bigoplus_{i \in I_n^1} P_n^i & \longrightarrow & \cdots & \longrightarrow & \bigoplus_{i \in I_1^1} P_1^i & \longrightarrow & \bigoplus_{i \in I_0^1} P_0^i & \longrightarrow & M_1 & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \bigoplus_{i \in I_n^2} P_n^i & \longrightarrow & \cdots & \longrightarrow & \bigoplus_{i \in I_1^2} P_1^i & \longrightarrow & \bigoplus_{i \in I_0^2} P_0^i & \longrightarrow & M_2 & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \bigoplus_{i \in I_n^2 - I_n^1} P_n^i & \longrightarrow & \cdots & \longrightarrow & \bigoplus_{i \in I_1^2 - I_1^1} P_1^i & \longrightarrow & \bigoplus_{i \in I_0^2 - I_0^1} P_0^i & \longrightarrow & M_2/M_1 & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
& & 0 & & 0 & & 0 & & 0 & & 0 & & 
\end{array}$$

where the first and third rows are exact sequences, and then so is the second since the class of exact complexes is closed under extensions. We have that  $M_1$  and  $M_2$  are  $n$ -projective modules such that  $M_1 \in \mathcal{P}_n^{\aleph_0}$ ,  $M_2/M_1 \in \mathcal{P}_n^{\aleph_0}$ . Now suppose that there is an ordinal  $\beta$  such that:

- (1)  $M_\alpha$  is an  $n$ -projective module, for every  $\alpha < \beta$ .
- (2)  $M_\alpha \subseteq M_{\alpha'}$  whenever  $\alpha \leq \alpha' < \beta$ .
- (3)  $M_{\alpha+1}/M_\alpha \in \mathcal{P}_n^{\aleph_0}$  whenever  $\alpha + 1 < \beta$ .
- (4)  $M_\gamma = \bigcup_{\alpha < \gamma} M_\alpha$  for every limit ordinal  $\gamma < \beta$ .

If  $\beta$  is a limit ordinal, then set  $M_\beta = \bigcup_{\alpha < \beta} M_\alpha$ . Otherwise there exists an ordinal  $\alpha < \beta$  such that  $\beta = \alpha + 1$ . In this case, construct  $M_{\alpha+1} \in \mathcal{P}_n$  from  $M_\alpha$  as we constructed  $M_2$  from  $M_1$ , such that  $M_{\alpha+1}/M_\alpha \in \mathcal{P}_n^{\aleph_0}$ . By transfinite induction, we obtain a  $\mathcal{P}_n^{\aleph_0}$ -filtration  $(M_\alpha : \alpha < \lambda)$  of  $M$ , for some ordinal  $\lambda$ , such that  $M_1 \supseteq N$  and  $M_1 \in \mathcal{P}_n^{\aleph_0}$ .

□

From now on,  $R$  shall be a noetherian ring. Now we are ready to prove the main result of this section.

**Theorem 3.2.** *Every chain complex  $X \in \text{dw}\widetilde{\mathcal{P}}_n$  has a  $\text{dw}\widetilde{\mathcal{P}}_n^{\aleph_0}$ -filtration.*

*Proof.* Let  $X \in \text{dw}\mathcal{P}_n$  and write

$$X = \cdots \longrightarrow X_{k+1} \xrightarrow{\partial_{k+1}} X_k \xrightarrow{\partial_k} X_{k-1} \longrightarrow \cdots.$$

For each  $k$  one has a projective resolution of  $X_k$  of length  $n$ :

$$0 \longrightarrow \bigoplus_{i \in I_n(k)} P_n^i(k) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1(k)} P_1^i(k) \longrightarrow \bigoplus_{i \in I_0(k)} P_0^i(k) \longrightarrow X_k \longrightarrow 0.$$

We shall construct a  $\text{dw}\widetilde{\mathcal{P}}_n^{\aleph_0}$ -filtration of  $X$  by using transfinite induction. For  $\alpha = 0$  set  $X^0 = 0$ . For  $\alpha = 1$ , choose  $m \in \mathbb{Z}$ . Let  $S$  be a countably generated submodule of  $X_m$ . By the previous lemma, there exists a submodule  $\mathcal{P}_n^{\aleph_0} \ni X_m^1 \subseteq X_m$  such that  $S \subseteq X_m^1$ . Note that  $X_m^1$  is also countably generated. Then  $\partial_m(X_m^1)$  is a countably generated submodule of  $X_{m-1}$ , and so there exists  $\mathcal{P}_n^{\aleph_0} \ni X_{m-1}^1 \subseteq X_{m-1}$  such that  $\partial_m(X_m^1) \subseteq X_{m-1}^1$ . Repeat the same procedure infinitely many times in order to obtain a subcomplex

$$X^1 = \cdots \longrightarrow X_{k+1}^1 \longrightarrow X_k^1 \longrightarrow X_{k-1}^1 \longrightarrow \cdots$$

of  $X$  such that  $X_k^1 \in \mathcal{P}_n^{\aleph_0}$  for every  $k \in \mathbb{Z}$  (we are setting  $X_k^1 = 0$  for every  $k > m$ ). Hence  $X^1 \in \text{dw}\widetilde{\mathcal{P}}_n^{\aleph_0}$ . Note from the proof of the previous lemma that the quotient  $X/X^1$  is in  $\text{dw}\widetilde{\mathcal{P}}_n$ . We have

$$X/X^1 = \cdots \longrightarrow X_{k+1}/X_{k+1}^1 \longrightarrow X_k/X_k^1 \longrightarrow X_{k-1}/X_{k-1}^1 \longrightarrow \cdots,$$

where for every  $k \leq m$  one has the following projective resolutions of length  $n$  for  $X_k^1$

and  $X_k/X_k^1$ :

$$\begin{aligned} 0 \longrightarrow \bigoplus_{i \in I_n^1(k)} P_n^i(k) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^1(k)} P_1^i(k) \longrightarrow \bigoplus_{i \in I_0^1(k)} P_0^i(k) \longrightarrow X_k^1 \longrightarrow 0, \\ 0 \longrightarrow \bigoplus_{i \in I_n(k) - I_n^1(k)} P_n^i(k) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_0(k) - I_0^1(k)} P_0^i(k) \longrightarrow X_k/X_k^1 \longrightarrow 0. \end{aligned}$$

Apply the same procedure above to the complex  $X/X^1$ , in order to get a subcomplex

$$X^2/X^1 = \cdots \longrightarrow X_{k+1}^2/X_{k+1}^1 \longrightarrow X_k^2/X_k^1 \longrightarrow X_{k-1}^2/X_{k-1}^1 \longrightarrow \cdots$$

of  $X/X^1$ , such that for each  $k \in \mathbb{Z}$  one has the following projective resolution of length  $n$  for  $X_k^2/X_k^1$ :

$$0 \longrightarrow \bigoplus_{i \in I_n^2 - I_n^1} P_n^i(k) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^2 - I_1^1} P_1^i(k) \longrightarrow \bigoplus_{i \in I_0^2 - I_0^1} P_0^i(k) \longrightarrow X_k^2/X_k^1 \longrightarrow 0,$$

where each  $I_j^2 - I_j^1 \subseteq I_j$  is countable. Now consider the complex

$$X^2 = \cdots \longrightarrow X_{k+1}^2 \longrightarrow X_k^2 \longrightarrow X_{k-1}^2 \longrightarrow \cdots.$$

As we did in the proof of the previous lemma, we have that

$$0 \longrightarrow \bigoplus_{i \in I_n^2(k)} P_n^i(k) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^2(k)} P_1^i(k) \longrightarrow \bigoplus_{i \in I_0^2(k)} P_0^i(k) \longrightarrow X_k^2 \longrightarrow 0$$

is an exact sequence. So  $X_k^2 \in \mathcal{P}_n$  for every  $k \in \mathbb{Z}$ , and hence  $X^2 \in \text{dw}\widetilde{\mathcal{P}}_n$ , with  $X^2/X^1 \in \text{dw}\widetilde{\mathcal{P}}_n^{\aleph_0}$ . The rest of the proof follows by transfinite induction, as in the end of the proof of the previous lemma.  $\square$

## 4 Exact degreewise $n$ -projective complexes

Consider the class of exact dw- $n$ -projective complexes  $\text{ex}\widetilde{\mathcal{P}}_n = \text{dw}\widetilde{\mathcal{P}}_n \cap \mathcal{E}$ , where  $\mathcal{E}$  denotes the class of exact complexes. The goal of this section is to prove that  $(\text{ex}\widetilde{\mathcal{P}}_n, (\text{ex}\widetilde{\mathcal{P}}_n)^\perp)$  is a complete cotorsion pair. This pair is a cotorsion pair by the following result by Gillespie:

**Proposition 4.1.** [8, Proposition 3.3] *Let  $(\mathcal{A}, \mathcal{B})$  be a cotorsion pair in an abelian category  $\mathcal{C}$  with enough projective and injective objects. If  $\mathcal{B}$  contains a cogenerator of finite injective dimension then  $(\text{ex}\widetilde{\mathcal{A}}, (\text{ex}\widetilde{\mathcal{A}})^\perp)$  is a cotorsion pair.*

Recall that a **cogenerator** in an abelian category  $\mathcal{C}$  is an object  $C$  such that for every nonzero object  $H$  there exists a nonzero morphism  $f : H \rightarrow C$ . For example,  $\text{Mod-}R$  has a an injective cogenerator given by the abelian group  $\text{Hom}(R, \mathbb{Q}/\mathbb{Z})$  of group homomorphisms and providing it with the scalar multiplication defined by

$$f \cdot r : R \rightarrow \mathbb{Q}/\mathbb{Z}, s \mapsto f(rs)$$

for  $f \in \text{Hom}(R, \mathbb{Q}/\mathbb{Z})$  and  $r \in R$  (see [3, Proposition 4.7.5] for details). Since  $\text{Hom}(R, \mathbb{Q}/\mathbb{Z}) \in \mathcal{P}_n^\perp$ , we have that  $(\text{ex}\mathcal{P}_n, (\text{ex}\mathcal{P}_n)^\perp)$  is a cotorsion pair.

Given a module  $M \in \mathcal{P}_n$ , consider a projective resolution of  $M$  of length  $n$ :

$$0 \rightarrow \bigoplus_{i \in I_n} P_n^i \rightarrow \bigoplus_{i \in I_{n-1}} P_{n-1}^i \rightarrow \cdots \rightarrow \bigoplus_{i \in I_1} P_1^i \rightarrow \bigoplus_{i \in I_0} P_0^i \rightarrow M \rightarrow 0 \quad (*).$$

We shall say that a projective resolution

$$0 \rightarrow \bigoplus_{i \in I'_n} P_n^i \rightarrow \bigoplus_{i \in I'_{n-1}} P_{n-1}^i \rightarrow \cdots \rightarrow \bigoplus_{i \in I'_1} P_1^i \rightarrow \bigoplus_{i \in I'_0} P_0^i \rightarrow N \rightarrow 0 \quad (**)$$

is a **nice subresolution** of  $(*)$  if  $I'_k \subseteq I_k$  for every  $0 \leq k \leq n$  and  $N \subseteq M$ .

From now on, fix an infinite cardinal  $\kappa$  such that  $\kappa \geq \text{Card}(R)$ . We shall say that a set  $S$  is **small** if  $\text{Card}(S) \leq \kappa$ . We shall also say that a chain complex  $X = (X_m)_{m \in \mathbb{Z}}$  is **small** if  $\text{Card}(X) \leq \kappa$ , where

$$\text{Card}(X) := \sum_{m \in \mathbb{Z}} \text{Card}(X_m).$$

So a complex  $X$  is small if and only if each term  $X_m$  is a small set. Note that if  $M$  is an  $n$ -projective module with a resolution given by  $(*)$ , then it is small if and only if  $\text{Card}(I_k) \leq \kappa$  for every  $0 \leq k \leq n$ . Let  $\mathcal{P}_n^{\leq \kappa}$  denote the set of  $n$ -projective modules with a small projective resolution. If we consider the resolutions  $(*)$  and  $(**)$  above, then note that  $(**)$  is a small and nice subresolution of  $(*)$  if each  $I'_k$  is a small subset of  $I_k$ . Consider the set

$$\widetilde{\text{ex}\mathcal{P}_n^{\leq \kappa}} = \{X \in \mathbf{Ch}(R\text{-}\mathbf{Mod}) : X \text{ is exact and } X_m \in \mathcal{P}_n^{\leq \kappa} \text{ for every } m \in \mathbb{Z}\}.$$

We shall prove that every exact dw- $n$ -projective complex has a  $\widetilde{\text{ex}\mathcal{P}_n^{\leq \kappa}}$ -filtration.

**Lemma 4.1.** *Let  $M \in \mathcal{P}_n$  with a projective resolution given by  $(*)$ . For every submodule  $N \subseteq M$  with  $\text{Card}(N) \leq \kappa$ , there exists a small and nice subresolution*

$$0 \longrightarrow \bigoplus_{i \in I'_n} P_n^i \longrightarrow \bigoplus_{i \in I'_{n-1}} P_{n-1}^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I'_1} P_1^i \longrightarrow \bigoplus_{i \in I'_0} P_0^i \longrightarrow N' \longrightarrow 0 \quad (***)$$

of  $(*)$  such that  $N \subseteq N'$ . Moreover, if  $N$  has an small and nice subresolution of  $M$ , then  $(***)$  can be constructed in such a way that it contains the given resolution of  $N$ .

*Proof.* Since  $f_0$  is surjective, for every  $x \in N$  choose  $y_x \in \bigoplus_{i \in I_0} P_0^i$  such that  $x = f_0(y_x)$ . Let  $Y = \{y_x : x \in N\}$ . Note that  $\langle Y \rangle$  is a small submodule of

$\bigoplus_{i \in I_0} P_0^i$ . So there exists a small subset  $I_0^0 \subseteq I_0$  such that  $\langle Y \rangle \subseteq \bigoplus_{i \in I_0^0} P_0^i$ . We have  $f_0 \left( \bigoplus_{i \in I_0^0} P_0^i \right) \supseteq N$ . Now consider the submodule  $\text{Ker} \left( f_0|_{\bigoplus_{i \in I_0^0} P_0^i} \right)$  of  $f_0 \left( \bigoplus_{i \in I_0^0} P_0^i \right)$ , which is small since  $f_0 \left( \bigoplus_{i \in I_0^0} P_0^i \right)$  is. Then we can choose a small subset  $I_1^0 \subseteq I_1$  such that  $f_1 \left( \bigoplus_{i \in I_1^0} P_1^i \right) \supseteq \text{Ker} \left( f_0|_{\bigoplus_{i \in I_0^0} P_0^i} \right)$ . Repeat the same argument until find a small subset  $I_n^0 \subseteq I_n$  such that  $f_n \left( \bigoplus_{i \in I_n^0} P_n^i \right) \supseteq \text{Ker} \left( f_{n-1}|_{\bigoplus_{i \in I_{n-1}^0} P_{n-1}^i} \right)$ . Since  $f_n \left( \bigoplus_{i \in I_n^0} P_n^i \right)$  is a small submodule of  $\bigoplus_{i \in I_{n-1}} P_{n-1}^i$ , we can choose a small subset  $I_{n-1}^0 \subseteq I_{n-1}^1 \subseteq I_{n-1}$  such that  $f_n \left( \bigoplus_{i \in I_n^0} P_n^i \right) \subseteq \bigoplus_{i \in I_{n-1}^1} P_{n-1}^i$ . From this point just use the zig-zag procedure in order to get small subsets  $I'_k = \bigcup_{j \geq 0} I_k^j \subseteq I_k$  and an exact sequence

$$0 \longrightarrow \bigoplus_{i \in I'_n} P_n^i \longrightarrow \bigoplus_{i \in I'_{n-1}} P_{n-1}^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I'_1} P_1^i \longrightarrow \bigoplus_{i \in I'_0} P_0^i \longrightarrow N' \longrightarrow 0$$

where  $N' := \text{CoKer} \left( \bigoplus_{i \in I'_1} P_1^i \longrightarrow \bigoplus_{i \in I'_0} P_0^i \right)$  and  $N \subseteq N' \subseteq M$ .

Now suppose that  $N$  has a small and nice subresolution

$$0 \longrightarrow \bigoplus_{i \in I'_n{}^N} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I'_1{}^N} P_1^i \longrightarrow \bigoplus_{i \in I'_0{}^N} P_0^i \longrightarrow N \longrightarrow 0$$

of  $(*)$ . Take the quotient of  $(*)$  by this resolution and get

$$0 \longrightarrow \bigoplus_{i \in I_n - I_n^N} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1 - I_1^N} P_1^i \longrightarrow \bigoplus_{i \in I_0 - I_0^N} P_0^i \longrightarrow M/N \longrightarrow 0.$$

Repeat the argument above using this sequence and the small submodule  $\langle z + N \rangle$ , where  $z \notin N$ . Then we get a projective subresolution of the previous one:

$$0 \longrightarrow \bigoplus_{i \in I'_n - I_n^N} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I'_1 - I_1^N} P_1^i \longrightarrow \bigoplus_{i \in I'_0 - I_0^N} P_0^i \longrightarrow N'/N \longrightarrow 0$$

where each set  $I'_k - I_k^N$  is a small set. As we did in the proof of Lemma 3.2, we have that

$$0 \longrightarrow \bigoplus_{i \in I'_n} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I'_1} P_1^i \longrightarrow \bigoplus_{i \in I'_0} P_0^i \longrightarrow N' \longrightarrow 0$$

is small and nice subresolution of  $(*)$ , and that contains the resolution of  $N$  as a nice subresolution.

□

**Lemma 4.2.** *Let  $X \in \widetilde{\text{dw}\mathcal{P}_n}$  and let  $Y$  be a bounded above subcomplex of  $X$  such that  $\text{Card}(Y_k) \leq \kappa$  for every  $k \in \mathbb{Z}$ . Then there exists a (bounded above) subcomplex  $Y'$  of  $X$  such that  $Y \subseteq Y'$  and  $Y' \in \widetilde{\text{dw}\mathcal{P}_n^{\leq \kappa}}$ .*

*Proof.* We are given the following commutative diagram

$$\begin{array}{ccccccc} Y = & \cdots & \longrightarrow & 0 & \longrightarrow & Y_m & \xrightarrow{\partial_m} & Y_{m-1} & \longrightarrow & \cdots \\ & & & \downarrow & & \downarrow & & \downarrow & & \\ X = & \cdots & \longrightarrow & X_{m+1} & \xrightarrow{\partial_{m+1}} & X_m & \xrightarrow{\partial_m} & X_{m-1} & \longrightarrow & \cdots \end{array}$$

Since  $X_m$  is an  $n$ -projective module, we have a projective resolution

$$0 \longrightarrow \bigoplus_{i \in I_n(m)} P_n^i(m) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1(m)} P_1^i(m) \longrightarrow \bigoplus_{i \in I_0(m)} P_0^i(m) \longrightarrow X_m \longrightarrow 0.$$

By the previous lemma, there exists a submodule  $Y'_m$  of  $X_m$  containing  $Y_m$ , along with a small and nice subresolution

$$0 \longrightarrow \bigoplus_{i \in I'_n(m)} P_n^i(m) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I'_1(m)} P_1^i(m) \longrightarrow \bigoplus_{i \in I'_0(m)} P_0^i(m) \longrightarrow Y'_m \longrightarrow 0.$$

Note that  $\text{Card}(\partial_m(Y'_m) + Y_{m-1}) \leq \kappa$  and  $Y_{m-1} \subseteq \partial_m(Y'_m) + Y_{m-1} \subseteq X_{m-1}$ . Now choose a submodule  $Y'_{m-1} \subseteq X_{m-1}$  such that  $\partial_m(Y'_m) + Y_{m-1} \subseteq Y'_{m-1}$  and  $Y'_{m-1}$  has a small and nice subresolution of a fixed resolution of  $X_{m-1}$ . Repeat this process

infinitely many times in order to obtain a complex

$$Y' = \cdots \longrightarrow 0 \longrightarrow Y'_m \longrightarrow Y'_{m-1} \longrightarrow \cdots$$

such that  $Y \subseteq Y' \subseteq X$  and  $Y' \in \text{dw}\widetilde{\mathcal{P}}_n^{\leq \kappa}$ .

□

**Theorem 4.1.** *Every  $X \in \text{ex}\widetilde{\mathcal{P}}_n$  has a  $\text{ex}\widetilde{\mathcal{P}}_n^{\leq \kappa}$ -filtration.*

*Proof.* Let  $X \in \text{ex}\widetilde{\mathcal{P}}_n$ . We construct a  $\text{ex}\widetilde{\mathcal{P}}_n^{\leq \kappa}$ -filtration of  $X$  using transfinite induction. For  $\alpha = 0$  set  $X^0 = 0$ . For the case  $\alpha = 1$  let  $m \in \mathbb{Z}$  be arbitrary and let  $T_1 \subseteq X_m$  be a small submodule of  $X_m$ . Then there exists a small submodule  $Y_m^1$  of  $X_m$  such that  $T_1 \subseteq Y_m^1$  and that  $Y_m^1$  has a small and nice projective subresolution of a given resolution of  $X_m$ . Note that  $\partial_m(Y_m^1)$  is a submodule of  $X_{m-1}$  with cardinality  $\leq \kappa$ , so there exists a submodule  $Y_{m-1}^1$  of  $X_{m-1}$  such that  $\partial_m(Y_m^1) \subseteq Y_{m-1}^1$  and that  $Y_{m-1}^1$  has a small and nice projective subresolution of a given resolution of  $X_{m-1}$ . Keep repeating this argument infinitely many times. We obtain a complex

$$Y^1 = \cdots \longrightarrow 0 \longrightarrow Y_m^1 \longrightarrow Y_{m-1}^1 \longrightarrow \cdots$$

which is a subcomplex of  $X$  and  $Y^1 \in \text{dw}\widetilde{\mathcal{P}}_n^{\leq \kappa}$ . Note that  $Y^1$  is not necessarily exact. We shall construct a complex  $X^1$  from  $Y^1$  such that  $X^1 \subseteq X$  and  $X^1 \in \text{ex}\widetilde{\mathcal{P}}_n^{\leq \kappa}$ . The rest of this proof uses an argument similar to the one used in [4, Theorem 4.6]. Fix any  $p \in \mathbb{Z}$ . Then  $\text{Card}(Y_p^1) \leq \kappa$  and so  $\text{Card}(Z_p Y^1) \leq \kappa$ . Since  $X$  is exact and  $\text{Card}(Z_p Y^1) \leq \kappa$ , there exists a submodule  $U \subseteq X_{p+1}$  with  $\text{Card}(U) \leq \kappa$  such that  $Z_p Y^1 \subseteq \partial_{p+1}(U)$ . Let  $C^1$  be a small subcomplex of  $X$  such that  $U \subseteq C_{p+1}$ ,  $C_j = 0$  for every  $j > p + 1$ , and that each  $C_j$  with  $j \leq p$  has a small and nice projective subresolution of a given resolution of  $X_j$ . Since  $Y^1 + C$  is a bounded above subcomplex of  $X$ , there exists a small subcomplex  $Y^2$  of  $X$  such that  $Y^1 + C \subseteq Y^2$  and that each  $Y_j^2$  has a small and nice projective subresolution of a given resolution of  $X_j$ . Note

that  $Z_p Y^1 \subseteq \partial_{p+1}(Y_{p+1}^2)$ . Construct  $Y^3$  from  $Y^2$  as we just constructed  $Y^2$  from  $Y^1$ , and so on, making sure to use the same  $p \in \mathbb{Z}$  at each step. Set  $X^1 = \bigcup_{j=1}^{\infty} Y^j \subseteq X$ . Note that  $X^1$  is exact at  $p$ . Repeat this argument to get exactness at any level. So we may assume that  $X^1$  is an exact complex. Every  $X_k^1$  has a small and nice projective subresolution of the given resolution of  $X_k$ . For every  $j$  one has a projective subresolution of the form

$$0 \longrightarrow \bigoplus_{i \in I_n^j(k)} P_n^i(k) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^j(k)} P_1^i(k) \longrightarrow \bigoplus_{i \in I_0^j(k)} P_0^i(k) \longrightarrow Y_k^j \longrightarrow 0,$$

where  $I_l^1(k) \subseteq I_l^2(k) \subseteq \cdots$  for every  $0 \leq l \leq n$ , by Lemma 4.1. If we take the union of all of the previous sequences, then we obtain the following exact sequence:

$$0 \longrightarrow \bigoplus_{i \in \bigcup_{j \geq 1} I_n^j(k)} P_n^i(k) \longrightarrow \cdots \longrightarrow \bigoplus_{i \in \bigcup_{j \geq 1} I_0^j(k)} P_0^i(k) \longrightarrow \bigcup_{j \geq 1} Y_k^j = X_k^1 \longrightarrow 0,$$

where  $\bigcup_{j \geq 1} I_l^j(k) \subseteq I_l(k)$  for every  $0 \leq l \leq n$ . Therefore,  $X^1 \in \widetilde{\text{ex}\mathcal{P}_n^{\leq \kappa}}$ . Now consider the quotient complex

$$X/X^1 = \cdots \longrightarrow X_{k+1}/X_{k+1}^1 \longrightarrow X_k/X_k^1 \longrightarrow X_{k-1}/X_{k-1}^1 \longrightarrow \cdots$$

Note that each  $X_k/X_k^1$  is  $n$ -projective and that  $X/X^1$  is exact. We apply the same procedure above to the complex  $X/X^1$  in order to get a complex  $X^2/X^1 \subseteq X/X^1$  such that  $X^2/X^1 \in \widetilde{\text{ex}\mathcal{P}_n^{\leq \kappa}}$ . Note that  $X^2$  is an exact complex since the class of exact complexes is closed under extensions, and so  $X^2 \in \widetilde{\text{ex}\mathcal{P}_n}$ . The rest of the proof follows by using transfinite induction. □

It follows by Proposition 3.2 and the Eklof and Trlifaj Theorem that  $(\widetilde{\text{ex}\mathcal{P}_n}, (\widetilde{\text{ex}\mathcal{P}_n})^\perp)$  is a complete cotorsion pair cogenerated by  $\widetilde{\text{ex}\mathcal{P}_n^{\leq \kappa}}$ .

## 5 The staircase zig-zag argument

In this section, we present another method to prove that the induced cotorsion pairs  $(\text{dw}\widetilde{\mathcal{P}}_n, (\text{dw}\widetilde{\mathcal{P}}_n)^\perp)$  and  $(\text{ex}\widetilde{\mathcal{P}}_n, (\text{ex}\widetilde{\mathcal{P}}_n)^\perp)$  are complete. This method also applies for the cotorsion pairs  $(\text{dw}\widetilde{\mathcal{F}}_n, (\text{dw}\widetilde{\mathcal{F}}_n)^\perp)$  and  $(\text{ex}\widetilde{\mathcal{F}}_n, (\text{ex}\widetilde{\mathcal{F}}_n)^\perp)$ . From now on, let  $\mathcal{A}$  denote either the class of projective modules or the class of flat modules, and let  $\mathcal{A}_n$  denote the class of all modules  $M$  having an  $\mathcal{A}$ -resolution of length  $n$ :

$$0 \longrightarrow A_n \longrightarrow \cdots \longrightarrow A_1 \longrightarrow A_0 \longrightarrow M \longrightarrow 0,$$

where  $A_k \in \mathcal{A}$  for every  $0 \leq k \leq n$ . Note that  $\mathcal{A}_0 = \mathcal{A}$ .

As we mentioned before, in [9] it is proven that  $(\mathcal{F}_n, \mathcal{F}_n^\perp)$  is a complete cotorsion pair. We give a simpler proof of this fact.

**Lemma 5.1.** *Let  $M \in \mathcal{F}_n$  with a flat resolution*

$$0 \longrightarrow F_n \xrightarrow{f_n} F_{n-1} \longrightarrow \cdots \longrightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} M \longrightarrow 0 \quad (1)$$

*and let  $N$  be a small submodule of  $M$ . Then there exists a flat subresolution*

$$0 \longrightarrow S'_n \longrightarrow \cdots \longrightarrow S'_1 \longrightarrow S'_0 \longrightarrow N' \longrightarrow 0$$

*of (1) such that  $S'_k$  is a small and pure submodule of  $F_k$ , for every  $0 \leq k \leq n$ , and such that  $N \subseteq N'$ . Moreover, if  $N$  has a subresolution of (1)*

$$0 \longrightarrow S_n \longrightarrow \cdots \longrightarrow S_1 \longrightarrow S_0 \longrightarrow N \longrightarrow 0$$

*where  $S_k$  is a small and pure submodule of  $F_k$ , for every  $0 \leq k \leq n$ , then the above resolution of  $N'$  can be constructed in such a way that it contains the resolution of  $N$ .*

*Proof.* First, note that for every flat module  $F$  and for every small submodule  $N \subseteq F$ , there exists a small and pure submodule  $S \subseteq F$  such that  $N \subseteq S$  (for a proof of this, see [6, Lemma 5.3.12]). For every  $x \in N$  there exists  $y_x \in F_0$  such that  $x = f_0(y_x)$ . Let  $Y$  be the set  $\{y_x : x \in N \text{ and } f_0(y_x) = x\}$  and consider the submodule  $\langle Y \rangle \subseteq F_0$ . Since  $\langle Y \rangle$  is small, there exists a small pure submodule  $S_0(1) \subseteq F_0$  such that  $\langle Y \rangle \subseteq S_0(1)$ . Note that  $f_0(S_0(1)) \supseteq N$ . Now consider  $\text{Ker}(f_0|_{S_0(1)})$  and let  $A$  be a set of preimages of  $\text{Ker}(f_0|_{S_0(1)})$  such that  $f_1(\langle A \rangle) \supseteq \text{Ker}(f_0|_{S_0(1)})$ . It is easy to see that  $\langle A \rangle$  is a small submodule of  $F_1$ , so we can embed it into a small pure submodule  $S_1(1) \subseteq F_1$ . Hence we have  $f_1(S_1(1)) \supseteq \text{Ker}(f_0|_{S_0(1)})$ . Now consider  $\text{Ker}(f_1|_{S_1(1)})$  and repeat the same process above in order to find a small pure submodule  $S_2(1) \subseteq F_2$  such that  $f_2(S_2(1)) \supseteq \text{Ker}(f_1|_{S_1(1)})$ . Keep doing this until find a small pure submodule  $S_n(1) \subseteq F_n$  such that  $f_n(S_n(1)) \supseteq \text{Ker}(f_{n-1}|_{S_{n-1}(1)})$ . Now  $f_n(S_n(1))$  is a small submodule of  $F_{n-1}$ , so there is a small pure submodule  $S_{n-1}(2) \subseteq F_{n-1}$  such that  $f_n(S_n(1)) \subseteq S_{n-1}(2)$ . Repeat this process until find a small pure submodule  $S_0(2) \subseteq F_0$  such that  $f_1(S_1(2)) \subseteq S_0(2)$ . If we now consider  $\text{Ker}(f_0|_{S_0(2)}) \subseteq F_0$ , we repeat the same argument above to find a small pure submodule  $S_1(3) \subseteq F_1$  such that  $f_1(S_1(3)) \supseteq \text{Ker}(f_0|_{S_0(2)})$ . Keep repeating this zig-zag procedure infinitely many times and set  $S_k = \bigcup_{i \geq 1} S_k(i)$ , for every  $0 \leq k \leq n$ . Note that each  $S_k$  is a pure submodule of  $F_k$ . By construction, we get an exact complex

$$0 \longrightarrow S_n \longrightarrow S_{n-1} \longrightarrow \cdots \longrightarrow S_1 \longrightarrow S_0 \longrightarrow Q \longrightarrow 0, \quad (2)$$

where  $Q = \text{CoKer}(f_1|_{S_1}) \subseteq M$ . If we take the quotient of (1) by (2), we get an exact complex

$$0 \longrightarrow F_n/S_n \longrightarrow F_{n-1}/S_{n-1} \longrightarrow \cdots \longrightarrow F_1/S_1 \longrightarrow F_0/S_0 \longrightarrow M/Q \longrightarrow 0.$$

Since each  $S_k$  is a pure submodule of  $F_k$ , we know that  $S_k$  and  $F_k/S_k$  are flat modules. Therefore,  $Q$  is a small  $n$ -flat submodule with  $N \subseteq Q$  such that  $M/Q$  is also  $n$ -flat. The rest of the proof follows as in Lemma 4.1.  $\square$

**Remark 5.1.** From Lemma 4.1 and Lemma 5.1, we have that for every  $A \in \mathcal{A}_n$  and for every small submodule  $0 \neq N \subseteq A$ , there exists a small submodule  $A' \subseteq A$  in  $\mathcal{A}$  such that  $N \subseteq A'$  and  $A/A' \in \mathcal{A}$ .

**Theorem 5.1.** Let  $X \in \text{ex}\widetilde{\mathcal{A}}_n$  and let  $x \in X$  (i.e.  $x \in X_m$  for some  $m \in \mathbb{Z}$ ). Then there exists a small complex  $Y \in \text{ex}\widetilde{\mathcal{A}}_n^{\leq \kappa}$  such that  $x \in Y$  and  $X/Y \in \text{ex}\widetilde{\mathcal{A}}_n$ .

The following proof is based on an argument given in [1, Proposition 4.1], where the authors prove that  $(\text{dw}\widetilde{\mathcal{F}}_0, (\text{dw}\widetilde{\mathcal{F}}_0)^\perp)$  and  $(\text{ex}\widetilde{\mathcal{F}}_0, (\text{ex}\widetilde{\mathcal{F}}_0)^\perp)$  are complete cotorsion pairs. We shall call this argument the **staircase zig-zag**.

*Proof of Theorem 5.1.* Assume without loss of generality that  $x \in X_0$ . Consider the submodule  $\langle x \rangle \subseteq X_0$ . Since  $X_0 \in \mathcal{A}$  and  $\langle x \rangle$  is small, we can embed  $\langle x \rangle$  into a submodule  $\mathcal{A}^{\leq \kappa} \ni Y_0^1 \subseteq X_0$  such that  $X_0/Y_0^1 \in \mathcal{A}$ . Since  $X$  is exact, we can construct a small and exact subcomplex

$$L^1 = \cdots L_2^1 \longrightarrow L_1^1 \longrightarrow Y_0^1 \longrightarrow \partial_0(Y_0^1) \longrightarrow 0 \longrightarrow \cdots .$$

Since  $\partial_0(Y_0^1)$  is small, there exists a submodule  $\mathcal{A}^{\leq \kappa} \ni Y_{-1}^2 \subseteq X_{-1}$  such that  $X_{-1}/Y_{-1}^2 \in \mathcal{A}$ . Now construct a small and exact subcomplex

$$L^2 = \cdots L_2^2 \longrightarrow L_1^2 \longrightarrow L_0^2 \longrightarrow Y_{-1}^2 \longrightarrow \partial_{-1}(Y_{-1}^2) \longrightarrow 0 \longrightarrow \cdots .$$

Note that it is possible to construct  $L^2$  containing  $L^1$ . Now embed  $L_0^2$  into a submodule  $\mathcal{A}^{\leq \kappa} \ni Y_0^3 \subseteq X_0$  such that  $X_0/Y_0^3 \in \mathcal{A}$ . Again, construct a small and exact subcomplex

$$L^3 = \cdots \longrightarrow L_2^3 \longrightarrow L_1^3 \longrightarrow Y_0^3 \longrightarrow Y_{-1}^2 + \partial_0(Y_0^3) \longrightarrow \partial_{-1}(Y_{-1}^2) \longrightarrow 0 \longrightarrow \cdots$$

containing  $L^2$ . Now let  $Y_1^4 \in \mathcal{A}^{\leq \kappa}$  be a submodule of  $X_1$  containing  $L_1^3$  such that  $X_1/Y_1^4 \in \mathcal{A}$ , and construct an exact and small complex

$$L^4 = \cdots \longrightarrow L_2^4 \longrightarrow Y_1^4 \longrightarrow Y_0^3 + \partial_1(Y_1^4) \longrightarrow Y_{-1}^2 + \partial_0(Y_0^3) \longrightarrow \partial_{-1}(Y_{-1}^2) \longrightarrow 0 \longrightarrow \cdots$$

containing  $L^3$ . Embed  $Y_0^3 + \partial_1(Y_1^4)$  into a submodule  $\mathcal{A}^{\leq \kappa} \ni Y_0^5 \subseteq X_0$  such that  $X_0/Y_0^5 \in \mathcal{A}$ . Construct an exact and small subcomplex

$$L^5 = \cdots \longrightarrow L_2^5 \longrightarrow L_1^5 \longrightarrow Y_0^5 \longrightarrow Y_{-1}^2 + \partial_0(Y_0^5) \longrightarrow \partial_{-1}(Y_{-1}^2) \longrightarrow 0 \longrightarrow \cdots$$


containing  $L^4$ . In a similar way, construct small and exact complexes

$$\begin{aligned} L^6 &= \cdots \longrightarrow L_1^6 \longrightarrow L_0^6 \longrightarrow Y_{-1}^6 \longrightarrow \partial_{-1}(Y_{-1}^6) \longrightarrow 0 \longrightarrow \cdots, \\ L^7 &= \cdots \longrightarrow L_1^7 \longrightarrow L_0^7 \longrightarrow L_{-1}^7 \longrightarrow Y_{-2}^7 \longrightarrow \partial_{-2}(Y_{-2}^7) \longrightarrow 0 \longrightarrow \cdots, \end{aligned}$$

such that  $Y_{-1}^6 \in \mathcal{A}$  is a small submodule of  $X_{-1}$  containing  $Y_{-1}^6 + \partial_0(Y_0^5)$ , and  $Y_{-2}^7 \in \mathcal{A}$  is a small submodule of  $X_{-2}$  containing  $\partial_{-1}(Y_{-1}^6)$ . We have the following commutative diagram of subcomplexes of  $X$ :

$$\begin{array}{cccccc}
0 & 0 & 0 & 0 & 0 & 0 \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^1 \longrightarrow & L_1^1 \longrightarrow & Y_0^1 \longrightarrow & \partial_0(Y_0^1) \longrightarrow & 0 \longrightarrow & 0 \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^2 \longrightarrow & L_1^2 \longrightarrow & L_0^2 \longrightarrow & Y_{-1}^2 \longrightarrow & \partial_{-1}(Y_{-1}^2) \longrightarrow & 0 \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^3 \longrightarrow & L_1^3 \longrightarrow & Y_0^3 \longrightarrow & \partial_0(Y_0^3) + Y_{-1}^2 \longrightarrow & \partial_{-1}(Y_{-1}^2) \longrightarrow & 0 \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^4 \longrightarrow & Y_1^4 \longrightarrow & Y_0^3 + \partial_1(Y_1^4) \longrightarrow & \partial_0(Y_0^3) + Y_{-1}^2 \longrightarrow & \partial_{-1}(Y_{-1}^2) \longrightarrow & 0 \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^5 \longrightarrow & L_1^5 \longrightarrow & Y_0^5 \longrightarrow & Y_{-1}^2 + \partial_0(Y_0^5) \longrightarrow & \partial_{-1}(Y_{-1}^2) \longrightarrow & 0 \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^6 \longrightarrow & L_1^6 \longrightarrow & L_0^6 \longrightarrow & Y_{-1}^6 \longrightarrow & \partial_{-1}(Y_{-1}^6) \longrightarrow & 0 \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^7 \longrightarrow & L_1^7 \longrightarrow & L_0^7 \longrightarrow & L_{-1}^7 \longrightarrow & Y_{-2}^7 \longrightarrow & \partial_{-2}(Y_{-2}^7) \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^8 \longrightarrow & L_1^8 \longrightarrow & L_0^8 \longrightarrow & Y_{-1}^8 \longrightarrow & \partial_{-1}(Y_{-1}^8) + Y_{-2}^7 \longrightarrow & \partial_{-2}(Y_{-2}^7) \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^9 \longrightarrow & L_1^9 \longrightarrow & Y_0^9 \longrightarrow & \partial_0(Y_0^9) + Y_{-1}^8 \longrightarrow & \partial_{-1}(Y_{-1}^8) + Y_{-2}^7 \longrightarrow & \partial_{-2}(Y_{-2}^7) \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
L_2^{10} \longrightarrow & Y_1^{10} \longrightarrow & \partial_1(Y_1^{10}) + Y_0^9 \longrightarrow & \partial_0(Y_0^9) + Y_{-1}^8 \longrightarrow & \partial_{-1}(Y_{-1}^8) + Y_{-2}^7 \longrightarrow & \partial_{-2}(Y_{-2}^7) \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
Y_2^{11} \longrightarrow & \partial_2(Y_2^{11}) + Y_1^{10} \longrightarrow & \partial_1(Y_1^{10}) + Y_0^9 \longrightarrow & \partial_0(Y_0^9) + Y_{-1}^8 \longrightarrow & \partial_{-1}(Y_{-1}^8) + Y_{-2}^7 \longrightarrow & \partial_{-2}(Y_{-2}^7) \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots
\end{array}$$

where the  $k+1$ -th complex can be constructed in such a way that it contains the  $k$ -th complex. Note that the submodules  $S_i^k$  appear according to the following pattern:



Let  $Y = \bigcup_{n \geq 1} L^n$ , where  $Y_i = \bigcup_{n \geq 1} (L^n)_i$ . It is clear that  $Y$  is an exact complex. We check that  $Y$  is also a  $\text{dw}\widetilde{\mathcal{A}}_n$ . For example, consider

$$Y_0 = Y_0^1 \cup L_0^2 \cup Y_0^3 \cup (Y_0^3 + \partial_1(Y_1^4)) \cup Y_0^5 \cup \dots = Y_0^1 \cup Y_0^3 \cup Y_0^5 \cup \dots.$$

It is clear that  $Y_0$  is small. At this point, we split the proof in two cases:

- (1)  $\mathcal{A} = \mathcal{P}_0$ : Consider a projective resolution of  $X_0$  of length  $n$ , say

$$0 \longrightarrow \bigoplus_{i \in I_n} P_n^i \longrightarrow \dots \longrightarrow \bigoplus_{i \in I_1} P_1^i \longrightarrow \bigoplus_{i \in I_0} P_0^i \longrightarrow X_0 \longrightarrow 0 \quad (1),$$

where each direct sum is a direct sum of countably generated projective modules.

By Lemma 4.1, we can construct  $Y_0^1$  containing  $\langle x \rangle$  with a subresolution

$$0 \longrightarrow \bigoplus_{i \in I_n^1} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^1} P_1^i \longrightarrow \bigoplus_{i \in I_0^1} P_0^i \longrightarrow Y_0^1 \longrightarrow 0 \quad (2),$$

where each  $I_k^1$  is a small subset of  $I_k$ . Note that the quotient of (1) by (2) yields a projective resolution of  $X_0/Y_0^1$  of length  $n$ , so  $X_0/Y_0^1 \in \mathcal{P}_n$ . Using Lemma 4.1 again, we can construct a subresolution

$$0 \longrightarrow \bigoplus_{i \in I_n^3} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^3} P_1^i \longrightarrow \bigoplus_{i \in I_0^3} P_0^i \longrightarrow Y_0^3 \longrightarrow 0 \quad (3)$$

containing (2) such that  $X_0/Y_0^3 \in \mathcal{P}_n$ . We keep applying Lemma 4.1 to get an ascending chain of subresolutions of (1):

$$\begin{aligned} 0 &\longrightarrow \bigoplus_{i \in I_n^1} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^1} P_1^i \longrightarrow \bigoplus_{i \in I_0^1} P_0^i \longrightarrow Y_0^1 \longrightarrow 0 \\ 0 &\longrightarrow \bigoplus_{i \in I_n^3} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^3} P_1^i \longrightarrow \bigoplus_{i \in I_0^3} P_0^i \longrightarrow Y_0^3 \longrightarrow 0 \\ 0 &\longrightarrow \bigoplus_{i \in I_n^5} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in I_1^5} P_1^i \longrightarrow \bigoplus_{i \in I_0^5} P_0^i \longrightarrow Y_0^5 \longrightarrow 0 \\ &\vdots \end{aligned}$$

Now we take the union of this ascending chain and get an exact complex

$$\begin{aligned} 0 &\longrightarrow \bigcup_j \bigoplus_{i \in I_n^j} P_n^i \longrightarrow \cdots \longrightarrow \bigcup_j \bigoplus_{i \in I_1^j} P_1^i \longrightarrow \bigcup_j \bigoplus_{i \in I_0^j} P_0^i \longrightarrow \bigcup_j Y_0^j \longrightarrow 0 \\ &= \\ 0 &\longrightarrow \bigoplus_{i \in \bigcup_j I_n^j} P_n^i \longrightarrow \cdots \longrightarrow \bigoplus_{i \in \bigcup_j I_1^j} P_1^i \longrightarrow \bigoplus_{i \in \bigcup_j I_0^j} P_0^i \longrightarrow Y_0 \longrightarrow 0 \quad (4) \end{aligned}$$

Since each  $\bigcup_j I_k^j$  is a small subset of  $I_k$ , we have that the previous sequence is a  $\mathcal{P}_0^{\leq \kappa}$ -subresolution of (1). Note also that the quotient of (1) by (4) yields a projective resolution of  $X_0/Y_0$  of length  $n$ . Then  $Y_0 \in \mathcal{P}_n^{\leq \kappa}$ . In a similar way, we can show that  $Y_m \in \mathcal{P}_n^{\leq \kappa}$  and  $X_m/Y_m \in \mathcal{P}_n$ , for every  $m \in \mathbb{Z}$ . Hence,  $Y \in \widetilde{\text{ex}\mathcal{P}_n^{\leq \kappa}}$ . Note also that the quotient of exact complexes is exact, so we also have  $X/Y \in \widetilde{\text{ex}\mathcal{P}_n^{\leq \kappa}}$ .

(2)  $\mathcal{A} = \mathcal{F}_0$ : Consider a flat resolution of  $X_0$  of length  $n$ , say

$$0 \longrightarrow F_n \longrightarrow \cdots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow X_0 \longrightarrow 0 \quad (1').$$

By Lemma 5.1, we can construct a subresolution

$$0 \longrightarrow S_n^1 \longrightarrow \cdots \longrightarrow S_1^1 \longrightarrow S_0^1 \longrightarrow Y_0^1 \longrightarrow 0,$$

where  $\langle x \rangle \subseteq Y_0^1$ , and each  $S_k^1$  is a small and pure submodule of  $F_k$ . As we did in the previous case, applying Lemma 5.1 infinitely many times, we can get an ascending chain of subresolutions

$$\begin{aligned} 0 &\longrightarrow S_n^1 \longrightarrow \cdots \longrightarrow S_1^1 \longrightarrow S_0^1 \longrightarrow Y_0^1 \longrightarrow 0 \\ 0 &\longrightarrow S_n^3 \longrightarrow \cdots \longrightarrow S_1^3 \longrightarrow S_0^3 \longrightarrow Y_0^3 \longrightarrow 0 \\ 0 &\longrightarrow S_n^5 \longrightarrow \cdots \longrightarrow S_1^5 \longrightarrow S_0^5 \longrightarrow Y_0^5 \longrightarrow 0 \\ &\vdots \end{aligned}$$

Taking the union of these subresolutions yields an exact sequence

$$0 \longrightarrow \bigcup_j S_n^j \longrightarrow \cdots \longrightarrow \bigcup_j S_1^j \longrightarrow \bigcup_j S_0^j \longrightarrow Y_0 \longrightarrow 0 \quad (2'),$$

where each  $\bigcup_j S_k^j$  is a small and pure submodule of  $F_k$ , and so it is flat and the

quotient  $F_k/\bigcup_j S_k^j$  is also flat. Then we have that  $Y_0 \in \mathcal{F}_n^{\leq \kappa}$  and  $X_0/Y_0 \in \mathcal{F}_n$  (take the quotient of (1') by (2')). In a similar way, we can show that  $Y_m \in \mathcal{F}_n^{\leq \kappa}$  and  $X_m/Y_m \in \mathcal{F}_n$ , for every  $m \in \mathcal{Z}$ . It follows  $Y \in \text{ex}\widetilde{\mathcal{F}_n^{\leq \kappa}}$  and  $X/Y \in \text{ex}\widetilde{\mathcal{F}_n}$ .

In both cases, we have constructed a subcomplex  $Y \subseteq X$  containing  $x$  such that  $Y \in \text{ex}\widetilde{\mathcal{A}_n^{\leq \kappa}}$  and  $X/Y \in \text{ex}\widetilde{\mathcal{A}_n}$ .

□

**Remark 5.2.** *Note that using the staircase zig-zag method, there is no need to assume that  $R$  is right noetherian.*

## 6 dw- $n$ -projective and dw- $n$ -flat model structures

In this section we obtain two model structures on  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$ , one from the two complete cotorsion pairs  $(\text{dw}\widetilde{\mathcal{P}_n}, (\text{dw}\widetilde{\mathcal{P}_n})^\perp)$  and  $(\text{ex}\widetilde{\mathcal{P}_n}, (\text{ex}\widetilde{\mathcal{P}_n})^\perp)$ , and the other one from  $(\text{dw}\widetilde{\mathcal{F}_n}, (\text{dw}\widetilde{\mathcal{F}_n})^\perp)$  and  $(\text{ex}\widetilde{\mathcal{F}_n}, (\text{ex}\widetilde{\mathcal{F}_n})^\perp)$ .

Recall that a subcategory  $\mathcal{D}$  of an abelian category  $\mathcal{C}$  is said to be **thick** if the following two conditions are satisfied:

- (a)  $\mathcal{D}$  is **closed under retracts**, i.e., given a sequence

$$D' \xrightarrow{f} D \xrightarrow{g} D'$$

with  $g \circ f = \text{id}_{D'}$  and  $D \in \mathcal{D}$ , then  $D' \in \mathcal{B}$ .

- (b) If two out of three of the terms in a short exact sequence

$$0 \longrightarrow D'' \longrightarrow D \longrightarrow D' \longrightarrow 0$$

are in  $\mathcal{D}$ , then so is the third.

For example, the class  $\mathcal{E}$  of exact complexes is thick.

The following theorem by Hovey describes how to get an abelian model structure from two complete cotorsion pairs.

**Theorem 6.1.** [10, Theorem 2.2] *Let  $\mathcal{C}$  be a bicomplete abelian category with enough projective and injective objects, and let  $(\mathcal{A}, \mathcal{B} \cap \mathcal{E})$  and  $(\mathcal{A} \cap \mathcal{E}, \mathcal{B})$  be two complete cotorsion pairs in  $\mathcal{C}$  such that the class  $\mathcal{E}$  is thick. Then there is a unique abelian model structure on  $\mathcal{C}$  such that  $\mathcal{A}$  is the class of cofibrant objects,  $\mathcal{B}$  is the class of fibrant objects, and  $\mathcal{E}$  is the class of trivial objects.*

Cotorsion pairs of the form  $(\mathcal{A}, \mathcal{B} \cap \mathcal{E})$  and  $(\mathcal{A} \cap \mathcal{E}, \mathcal{B})$  are called **compatible** by Gillespie in [8]. We shall use the following result from [4] to show that  $(\mathrm{dw}\widetilde{\mathcal{P}}_n, (\mathrm{dw}\widetilde{\mathcal{P}}_n)^\perp)$  and  $(\mathrm{ex}\widetilde{\mathcal{P}}_n, (\mathrm{ex}\widetilde{\mathcal{P}}_n)^\perp)$  are compatible cotorsion pairs.

**Lemma 6.1.** [4, Lemma 5.3] *If  $(\mathcal{C}, \mathcal{D}')$  is a cotorsion pair and  $(\mathcal{U}, \mathcal{V})$  is a complete and hereditary cotorsion pair in  $\mathbf{Ch}(R\text{--}\mathbf{Mod})$  (i.e.  $\mathcal{V}$  contains the injective complexes and it is closed under extensions and under taking cokernels of monomorphisms), and if  $\mathcal{U} \subseteq \mathcal{C}$  then when  $(\mathcal{C} \cap \mathcal{V})^\perp = \mathcal{D}$ , we have  $\mathcal{D}' = \mathcal{D} \cap \mathcal{V}$ .*

In the previous lemma, put  $(\mathcal{C}, \mathcal{D}') = (\mathrm{dw}\widetilde{\mathcal{A}}_n, (\mathrm{dw}\widetilde{\mathcal{A}}_n)^\perp)$  and  $(\mathcal{U}, \mathcal{V}) = ({}^\perp\mathcal{E}, \mathcal{E})$ . In [4, Lemma 5.1] it is proven that  $({}^\perp\mathcal{E}, \mathcal{E})$  is cogenerated by a set, so it is complete. In [7], it is proven that  ${}^\perp\mathcal{E}$  is the class of dg-projective complexes, where a complex  $X$  is dg-projective if  $X_m \in \mathcal{P}_0$  for every  $m \in \mathbb{Z}$  and every map  $X \rightarrow Y$  is nullhomotopic whenever  $Y$  is exact. Hence, it is clear that  ${}^\perp\mathcal{E} \subseteq \mathrm{dw}\widetilde{\mathcal{A}}_n$ , i.e.  $\mathcal{U} \subseteq \mathcal{C}$ . Setting  $\mathcal{D} = (\mathcal{C} \cap \mathcal{V})^\perp = (\mathrm{dw}\widetilde{\mathcal{A}}_n \cap \mathcal{E})^\perp$ , we have  $(\mathrm{dw}\widetilde{\mathcal{A}}_n)^\perp = (\mathrm{dw}\widetilde{\mathcal{A}}_n \cap \mathcal{E})^\perp \cap \mathcal{E}$ . So we obtain

$$\begin{aligned}
(\mathrm{dw}\widetilde{\mathcal{A}}_n, (\mathrm{dw}\widetilde{\mathcal{A}}_n)^\perp) &= (\mathrm{dw}\widetilde{\mathcal{A}}_n, (\mathrm{dw}\widetilde{\mathcal{A}}_n \cap \mathcal{E})^\perp \cap \mathcal{E}), \\
(\mathrm{ex}\widetilde{\mathcal{A}}_n, (\mathrm{ex}\widetilde{\mathcal{A}}_n)^\perp) &= (\mathrm{dw}\widetilde{\mathcal{A}}_n \cap \mathcal{E}, (\mathrm{dw}\widetilde{\mathcal{A}}_n \cap \mathcal{E})^\perp).
\end{aligned}$$

From Theorem 6.1 and the previous equalities, we have:

**Corollary 6.1.** *There exists a unique abelian model structure in  $\mathbf{Ch}(R\text{-}\mathbf{Mod})$  such that  $\mathrm{dw}\widetilde{\mathcal{A}}_n$  is the class of cofibrant objects,  $(\mathrm{ex}\widetilde{\mathcal{A}}_n)^\perp$  is the class of fibrant objects, and  $\mathcal{E}$  is the class of trivial objects. In the case  $\mathcal{A} = \mathcal{P}_0$  we name this structure the **dw- $n$ -projective model structure**, and when  $\mathcal{A} = \mathcal{F}_0$  we name it the **dw- $n$ -flat model structure**.*

**Remark 6.1.** *As far as the author knows, the dw-projective model structure first appeared in [4], while the dw-flat model did in [8].*

## References

- [1] S. T. Aldrich, E. E. Enochs, J. R. García Rozas & L. Oyonarte. *Covers and Envelopes in Grothendieck Categories: Flat Covers of Complexes with Applications*. Journal of Algebra. Vol. 243. pp. 615-630. (2001).
- [2] S. T. Aldrich, E. E. Enochs, O. M. G. Jenda & L. Oyonarte. *Envelopes and Covers by Modules of Finite Injective and Projective Dimensions*. Journal of Algebra, Vol. 242, pp. 447-459. (2001).
- [3] F. Borceaux. *Handbook of Categorical Algebra 1. Basic Category Theory*. Encyclopedia of Mathematics and its Applications, Vol. 50. Cambridge University Press. (1994).

- [4] D. Bravo, E. E. Enochs, A. C. Iacob, O. M. G. Jenda & J. Rada. *Cotorsion pairs in  $C(R\text{-Mod})$* . To appear in Rocky Mountain Journal of Mathematics.
- [5] P. Eklof & J. Trlifaj. *How to make Ext vanish*. Bulletin of the London Mathematical Society, Vol. 33, No. 1, pp. 41-51. (2001).
- [6] E. E. Enochs & O. M. G. Jenda. *Relative Homological Algebra*. De Gruyter Expositions in Mathematics, Vol. 30. De Gruyter. Berlin. (2000).
- [7] E.E. Enochs, O.M.G. Jenda & J. Xu. *Orthogonality in the category of complexes*. Math. J. Okayama Univ. Vol. 38. pp. 25-46. (1996).
- [8] J. Gillespie. *Cotorsion Pairs and Degreewise Homological Model Structures*. Homology, Homotopy and Applications, Vol. 10, No. 1, pp. 283-304. (2008).
- [9] R. Göbel & J. Trlifaj. *Approximations and Endomorphism Algebras of Modules*. De Gruyter Expositions in Mathematics, Vol. 31. De Gruyter. Berlin. (2006).
- [10] M. Hovey. *Cotorsion pairs, model category structures and representation theory*.
- [11] I. Kaplansky. *Projective modules*. Annals of Mathematics. Vol. 68 , no. 2, pp. 372-377 (1958).